arXiv:1702.06328v1 [math.GN] 21 Feb 2017

On selective sequentially separability of space of Borel functions[☆]

Alexander V. Osipov

Krasovskii Institute of Mathematics and Mechanics, Ural Federal University, Ekaterinburg, Russia

Abstract

In paper we proved that:

• a space of Borel functions $B(X)$ on a set of reals X, with pointwise topology, to be selective sequentially separable if and only, if X has a property $S_1(B_\Gamma, B_\Gamma)$.

• there exists a consistent example of sequentially separable selectively separable space which is not selective sequentially separable. This is an answer to a question of A. Bella, M. Bonanzinga and M. Matveev.

• there is a consistent example of a compact T_2 sequentially separable space which is not selective sequentially separable. This is an answer to a question of A. Bella and C. Costantini.

• $\min\{\mathfrak{b}, \mathfrak{q}\} = \{\kappa : 2^\kappa$ is not selective sequentially separable$\}$. This is an answer to a question of A. Bella, M. Bonanzinga and M. Matveev.

Keywords: $S_1(\mathcal{D}, \mathcal{D})$, $S_1(\mathcal{S}, \mathcal{S})$, $S_{fin}(\mathcal{S}, \mathcal{S})$, function spaces, selection principles, Baire function, σ-set, $S_1(B_\Omega, B_\Omega)$, $S_1(B_\Gamma, B_\Gamma)$, $S_1(B_\Omega, B_\Gamma)$, sequentially separable, selectively separable, selective sequentially separable
2000 MSC: 37F20, 26A03, 03E75, 54C35

1. Introduction

In [7], Osipov and Pytkeev gave necessary and sufficient conditions for the space $B_1(X)$ of first Baire class functions on a Tychonoff space X, with pointwise topology, to be (strongly) sequentially separable. In this paper we

[☆]The research has been supported by .
Email address: OAB@list.ru (Alexander V. Osipov)

consider some properties of a space $B(X)$ of Borel functions on a set of reals X, with pointwise topology, that are stronger than separability.

Many topological properties are defined or characterized in terms of the following classical selection principles. Let $\mathcal{A}$ and $\mathcal{B}$ be sets consisting of families of subsets of an infinite set X. Then:

$S_1(\mathcal{A},\mathcal{B})$ is the selection hypothesis: for each sequence $\{A_n : n \in \omega\}$ of elements of $\mathcal{A}$ there is a sequence $\{b_n : n \in \omega\}$ such that for each n, $b_n \in A_n$, and $\{b_n : n \in \omega\}$ is an element of $\mathcal{B}$.

$S_{fin}(\mathcal{A},\mathcal{B})$ is the selection hypothesis: for each sequence $\{A_n : n \in \omega\}$ of elements of $\mathcal{A}$ there is a sequence $\{B_n : n \in \omega\}$ of finite sets such that for each n, $B_n \subseteq A_n$, and $\bigcup_{n\in\omega} B_n \in \mathcal{B}$.

$U_{fin}(\mathcal{A},\mathcal{B})$ is the selection hypothesis: whenever $\mathcal{U}_1, \mathcal{U}_2, ... \in \mathcal{A}$ and none contains a finite subcover, there are finite sets $\mathcal{F}_n \subseteq \mathcal{U}_n$, $n \in \omega$, such that $\{\bigcup \mathcal{F}_n : n \in \omega\} \in \mathcal{B}$.

2. Main definitions and notation

An open cover $\mathcal{U}$ of a space X is:

• an *ω-cover* if X does not belong to $\mathcal{U}$ and every finite subset of X is contained in a member of $\mathcal{U}$.

• a *γ-cover* if it is infinite and each $x \in X$ belongs to all but finitely many elements of $\mathcal{U}$.

For a topological space X we denote:

• Ω — the family of countable open ω-covers of X;

• Γ — the family of countable open γ-covers of X;

• B_Ω — the family of countable Borel ω-covers of X;

• B_Γ — the family of countable Borel γ-covers of X;

• F_Γ — the family of countable closed γ-covers of X;

• $\mathcal{D}$ — the family of countable dense subsets of X;

• $\mathcal{S}$ — the family of countable sequentially dense subsets of X.

We will be denoted by

• $C_p(X)$ a set of all real-valued continuous functions $C(X)$ defined on a space X, with pointwise topology.

• $B_1(X)$ a set of all first Baire class functions $B_1(X)$ i.e., pointwise limits of continuous functions, defined on a space X, with pointwise topology.

• $B(X)$ a set of all Borel functions, defined on a space X, with pointwise topology.

If X is a space and $A \subseteq X$, then the sequential closure of A, denoted by $[A]_{seq}$, is the set of all limits of sequences from A. A set $D \subseteq X$ is said to be sequentially dense if $X = [D]_{seq}$. If D is a countable sequentially dense subset of X then X call sequentially separable space.

Call X strongly sequentially separable if X is separable and every countable dense subset of X is sequentially dense.

Recall that the cardinal $\mathfrak{p}$ is the smallest cardinal so that there is a collection of $\mathfrak{p}$ many subsets of the natural numbers with the strong finite intersection property but no infinite pseudo-intersection. Note that $\omega_1 \leq \mathfrak{p} \leq \mathfrak{c}$.

For $f, g \in \mathbb{N}^{\mathbb{N}}$, let $f \leq^* g$ if $f(n) \leq g(n)$ for all but finitely many n. $\mathfrak{b}$ is the minimal cardinality of a $\leq^*$-unbounded subset of $\mathbb{N}^{\mathbb{N}}$. A set $B \subset [\mathbb{N}]^{\infty}$ is unbounded if the set of all increasing enumerations of elements of B is unbounded in $\mathbb{N}^{\mathbb{N}}$, with respect to $\leq^*$. It follows that $|B| \geq \mathfrak{b}$. A subset S of the real line is called a Q-set if each one of its subsets is a G_δ. The cardinal $\mathfrak{q}$ is the smallest cardinal so that for any $\kappa < \mathfrak{q}$ there is a Q-set of size κ. (See [6] for more on small cardinals including $\mathfrak{p}$).

A space X is selectively separable (or M-separable, see for example [3]) if for every sequence $\{D_n : n \in \omega\}$ of dense subsets of X one can pick finite $F_n \subset D_n$, $n \in \omega$, so that $\bigcup\{F_n : n \in \omega\}$ is dense in X.

In [3] the authors started to investigate a selective version of sequential separability.

A space X is selectively sequentially separable (or M-sequentially separable, see for example [3]) if for every sequence $\{D_n : n \in \omega\}$ of sequentially dense subsets of X, one can pick finite $F_n \subset D_n$, $n \in \omega$, so that $\bigcup\{F_n : n \in \omega\}$ is sequentially dense in X.

In Scheeper's terminology [11] selectively separability equivalently to the selection principle $S_{fin}(\mathcal{D}, \mathcal{D})$, and selective sequentially separability equivalently to the $S_{fin}(\mathcal{S}, \mathcal{S})$.

3. Properties of a space of Borel functions

Theorem 3.1. *For a set of reals X, the following statements are equivalent:*

1. *$B(X)$ is sequentially separable and $B(X) \in S_1(\mathcal{S}, \mathcal{S})$;*
2. *$X \in S_1(B_\Gamma, B_\Gamma)$;*
3. *$B(X)$ is sequentially separable and $B(X) \in S_{fin}(\mathcal{S}, \mathcal{S})$;*
4. *$X \in S_{fin}(B_\Gamma, B_\Gamma)$;*
5. *$B_1(X) \in S_1(\mathcal{S}, \mathcal{S})$;*

6. $X \in S_1(F_\Gamma, F_\Gamma)$;
7. $B_1(X) \in S_{fin}(\mathcal{S}, \mathcal{S})$.

Proof. It is obviously that (1) $\Rightarrow$ (3).

(2) $\Leftrightarrow$ (4). By Theorem 1 in [10], $U_{fin}(B_\Gamma, B_\Gamma) = S_1(B_\Gamma, B_\Gamma) = S_{fin}(B_\Gamma, B_\Gamma)$.

(3) $\Rightarrow$ (2). Let $\{\mathcal{F}_i\} \subset B_\Gamma$ and $\mathcal{S} = \{h_m\}_{m\in\omega}$ be a countable sequentially dense subset of $B(X)$. For each $i \in \omega$ we consider a countable sequentially dense subset $\mathcal{S}_i$ of $B(X)$ and $\mathcal{F}_i = \{F_i^m\}_{m\in\omega}$ where

$\mathcal{S}_i = \{f_i^m\} := \{f_i^m \in B(X) : f_i^m \restriction F_i^m = h_m$ and $f_i^m \restriction (X \setminus F_i^m) = 1$ for $m \in \omega\}$.

Since $\mathcal{F}_i = \{F_i^m\}_{m\in\omega}$ is a Borel γ-cover of X and $\mathcal{S}$ is a countable sequentially dense subset of $B(X)$, we have that $\mathcal{S}_i$ is a countable sequentially dense subset of $B(X)$ for each $i \in \omega$. Really, let $h \in B(X)$, there is a sequence $\{h_s\}_{s\in\omega} \subset \mathcal{S}$ such that $\{h_s\}_{s\in\omega}$ converge to h. We claim that $\{f_i^s\}_{s\in\omega}$ converge to h. Let $K = \{x_1, ..., x_k\}$ be a finite subset of X, $\epsilon > 0$ and $W =< h, K, \epsilon >:= \{g \in B(X) : |g(x_j) - h(x_j)| < \epsilon$ for $j = 1, ..., k\}$ be a base neighborhood of h, then there is a number m_0 such that $K \subset F_i^m$ for $m > m_0$ and $h_s \in W$ for $s > m_0$. Since $f_i^s \restriction K = h_s \restriction K$ for each $s > m_0$, $f_i^s \in W$ for each $s > m_0$. It follows that a sequence $\{f_i^s\}_{s\in\omega}$ converge to h.

Since $B(X) \in S_{fin}(\mathcal{S}, \mathcal{S})$, there is a sequence $\{F_i = \{f_i^{m_1}, ..., f_i^{m_{s(i)}}\} : i \in \omega\}$ such that for each i, $F_i \subset \mathcal{S}_i$, and $\bigcup_{i\in\omega} F_i$ is a countable sequentially dense subset of $B(X)$.

For $\mathbf{0} \in B(X)$ there is a sequence $\{f_{i_j}^{m_{j(i)}}\}_{j\in\omega} \subset \bigcup_{i\in\omega} F_i$ such that $\{f_{i_j}^{m(i)_j}\}_{j\in\omega}$ converge to $\mathbf{0}$. Consider a sequence $\{F_{i_j}^{m_{j(i)}} : j \in \omega\}$.

(1) $F_{i_j}^{m_{j(i)}} \in \mathcal{F}_{i_j}$.

(2) $\{F_{i_j}^{m_{j(i)}} : j \in \omega\}$ is a γ-cover of X.

Really, let K be a finite subset of X and $U =< \mathbf{0}, K, \frac{1}{2} >$ be a base neighborhood of $\mathbf{0}$, then there is a number j_0 such that $f_{i_j}^{m_{j(i)}} \in U$ for any $j > j_0$. It follows that $K \subset F_{i_j}^{m_{j(i)}}$ for any $j > j_0$. We thus get $X \in U_{fin}(B_\Gamma, B_\Gamma)$, and, hence, by Theorem 1 in [10], $X \in S_1(B_\Gamma, B_\Gamma)$.

(2) $\Rightarrow$ (1). Let $\{S_i\} \subset \mathcal{S}$ and $S \in \mathcal{S}$. Consider a topology τ generated by the family $\mathcal{P} = \{f^{-1}(G) : G$ is an open set of $\mathbb{R}$ and $f \in S \cup \bigcup\limits_{i\in\omega} S_i\}$. A space $Y = (X, \tau)$ is a separable metrizable space because $P = S \cup \bigcup\limits_{i\in\omega} S_i$ is a countable dense subset of $B(X)$. Note that a function $f \in P$, considered as mapping from Y to $\mathbb{R}$, is a continuous function. Note also that the identity map φ from X on Y, is a Borel bijection. By Corollary 12 in [5], Y is a

QN-space and, hence, by Corollary 20 in [12], Y has a property $S_1(B_\Gamma, B_\Gamma)$. By Corollary 21 in [12], $B(Y)$ is an α_2 space and, hence, $B(Y)$ has property $S_1(\Gamma_z, \Gamma_z)$ for each $z \in B(Y)$.

We claim that there is a sequence $B = \{b_i : i \in \omega\}$ such that for each i, $b_i \in S_i \subset C(Y)$, and, $S \subset [B]_{seq}$, i.e. for each $d \in S \subset C(Y)$ there is a sequence $\{b_{i_k} : k \in \omega\} \subset B$ such that $\{b_{i_k} : k \in \omega\}$ converges to d.

Let $q : \omega \mapsto \omega \times \omega$ be a bijection. Then we enumerate $\{S_i\}$ as $\{S_{q(i)}\}$. For each $d_n \in S$ there are sequences $s_{n,m} \subset S_{n,m}$ such that $s_{n,m}$ converges to d_n for each $m \in \omega$. By property $S_1(\Gamma_z, \Gamma_z)$ of $B(Y)$, there is a sequence $\{b_{n,m} : m \in \omega\}$ such that for each m, $b_{n,m} \in s_{n,m}$, and, $\{b_{n,m}\}$ converges to d_n. Denote $B = \{b_{n,m} : n, m \in \omega\}$. By construction of B, we have that $S \subset [B]_{seq}$.

Since X is a σ-set (see [12]), $B_1(X) = B(X)$ and $\varphi(B(Y)) = \varphi(B_1(Y)) \subseteq B(X)$. Since S is a countable sequentially dense subset of $B(X)$, for any $g \in B(X)$ there is a sequence $\{g_n\}_{n\in\omega} \subset S$ such that $\{g_n\}_{n\in\omega}$ converge to g. But g we can consider as a mapping from Y into $\mathbb{R}$ and a sequence $\{g_n\}_{n\in\omega}$ as subset of $C(Y)$. It follows that $g \in B_1(Y)$. We get that $\varphi(B(Y)) = B(X)$.

We claim that $B \in \mathcal{S}$. Let $f \in B(Y)$ and $\{f_k : k \in \omega\} \subset S$ such that $\{f_k\}$ converge to f. For each $k \in \omega$ there is $\{f_k^n : n \in \omega\} \subset B$ such that $\{f_k^n\}$ converge to f_k. Since Y is a QN-space, by Theorem 16 in [5], there exists an unbounded $\beta \in \omega^\omega$ such that $\{f^n_{\beta(n)}\}$ converge to f on Y. It follows that $\{f^n_{\beta(n)}\}$ converge to f on X and $[B]_{seq} = B(X)$.

(5) $\Rightarrow$ (6). The Velichko's Theorem in [13] shows that a space $B_1(X)$ is sequentially separable for any separable metric space X.

Let $\{\mathcal{F}_i\} \subset F_\Gamma$ and $\mathcal{S} = \{h_m\}_{m\in\omega}$ be a countable sequentially dense subset of $B_1(X)$.

Similarly implication (3) $\Rightarrow$ (2) we get $X \in U_{fin}(F_\Gamma, F_\Gamma)$, and, hence, by Lemma 13 in [12], $X \in S_1(F_\Gamma, F_\Gamma)$.

(6) $\Rightarrow$ (5). By Corollary 20 in [12], $X \in S_1(B_\Gamma, B_\Gamma)$. Since X is a σ-set (see [12]), $B_1(X) = B(X)$ and, by implication (2) $\Rightarrow$ (1), we get $B_1(X) \in S_1(\mathcal{S}, \mathcal{S})$.

□

In [11] (Theorem 13) M. Scheeper proved the following result

Theorem 3.2. *(Scheeper) For X a separable metric space, the following are equivalent:*

1. *$C_p(X)$ has property $S_1(\mathcal{D}, \mathcal{D})$;*

2. *X has property $S_1(\Omega, \Omega)$.*

We claim the theorem for a space $B(X)$ of Borel functions.

Theorem 3.3. *For a set of reals X, the following are equivalent:*

1. $B(X) \in S_1(\mathcal{D}, \mathcal{D})$;
2. $X \in S_1(B_\Omega, B_\Omega)$.

Proof. (1) $\Rightarrow$ (2). Let X be a set of reals satisfying the hypotheses and β be a countable base of X. Consider a sequence $\{\mathcal{B}_i\}_{i\in\omega}$ of countable Borel ω-covers of X where $\mathcal{B}_i = \{W_i^j\}_{j\in\omega}$ for each $i \in \omega$.

Consider a topology τ generated by the family $\mathcal{P} = \{W_i^j \cap A_s : i, j \in \omega$ and $A_s \in \beta\} \bigcup \{(X \setminus W_i^j) \cap A_s : i, j \in \omega$ and $A_s \in \beta\}$.

Note that if χ_P is a characteristic function of P for each $P \in \mathcal{P}$, then a diagonal mapping $\varphi = \Delta_{P\in\mathcal{P}}\chi_P : X \mapsto 2^\omega$ is a Borel bijection. Let $Z = \varphi(X)$.

A space $Z = (X, \tau)$ is a separable metrizable space. Note that $\{\mathcal{B}_i\}$ is countable open ω-cover of Z for each $i \in \omega$. Since $B(\varphi(X)) = B(Z)$ is a dense subset of $B(X)$, then $B(Z)$ also has property $S_1(\mathcal{D}, \mathcal{D})$. Since $C_p(Z)$ is a dense subset of $B(Z)$, $C_p(Z)$ also has property $S_1(\mathcal{D}, \mathcal{D})$.

By Scheeper's Theorem 3.2, the space Z has property $S_1(\Omega, \Omega)$. It follows that there is a sequence $\{W_i^{j(i)}\}_{i\in\omega}$ such that $W_i^{j(i)} \in \mathcal{B}_i$ and $\{W_i^{j(i)}\}_{i\in\omega}$ is open ω-cover of Z. It follows that $\{W_i^{j(i)}\}_{i\in\omega}$ is Borel ω-cover of X.

(2) $\Rightarrow$ (1). Assume that X has property $S_1(B_\Omega, B_\Omega)$. Let $\{D_i\}_{i\in\omega}$ be a sequence countable dense subsets of $B(X)$. We claim that for any $f \in B(X)$ there is a sequence $\{f_k\} \subset B(X)$ such that $f_k \in D_k$ for each $k \in \omega$ and $f \in \overline{\{f_k : k \in \omega\}}$. Without loss of generality we can assume $f = \mathbf{0}$. Denote $W_i^k = \{x \in X : -\frac{1}{k} < f_i^k(x) < \frac{1}{k}\}$ for each $f_i^k \in D_k = \{f_i^k : i \in \omega\}$ and $k \in \omega$.

If for each $j \in \omega$ there is $k(j)$ such that $W_{i(j)}^{k(j)} = X$, then a sequence $f_{k(j)} = f_{i(j)}^{k(j)}$ uniform convergence to f and, hence, $f \in \overline{\{f_{k(j)} : j \in \omega\}}$.

We can assume that $W_i^k \neq X$ for any $k, i \in \omega$.

(a). $\{W_i^k\}_{i\in\omega}$ a sequence of Borel sets of X.

(b). For each $k \in \omega$, $\{W_i^k\}_{i\in\omega}$ is a ω-cover of X.

By (2), X has property $S_1(B_\Omega, B_\Omega)$, hence, there is a sequence $\{W_{i(k)}^k\}_{k\in\omega}$ such that $W_{i(k)}^k \in \{W_i^k\}_{i\in\omega}$ for each $k \in \omega$ and $\{W_{i(k)}^k\}_{k\in\omega}$ is a ω-cover of X.

Consider $\{f_{i(k)}^k\}$ and we claim that $f \in \overline{\{f_{i(k)}^k : k \in \omega\}}$. Let K be a finite subset of X, $\epsilon > 0$ and $U =< f, K, \epsilon >$ be a base neighborhood of f, then there is $k_0 \in \omega$ such that $\frac{1}{k_0} < \epsilon$ and $K \subset W_{i(k_0)}^{k_0}$. It follows that $f_{i(k_0)}^{k_0} \in U$.

Let $D = \{d_n : n \in \omega\}$ be a dense subspace of $B(X)$. Given a sequence $\{D_i\}_{i\in\omega}$ of dense subspace of $B(X)$, enumerate it as $\{D_{n,m} : n, m \in \omega\}$. For each $n \in \omega$, pick $d_{n,m} \in D_{n,m}$ so that $d_n \in \overline{\{d_{n,m} : m \in \omega\}}$. Then $\{d_{n,m} : m, n \in \omega\}$ is dense in $B(X)$.

□

Theorem 3.4. *For a set of reals X, the following are equivalent:*

1. $B(X) \in S_{fin}(\mathcal{D}, \mathcal{D})$;
2. $X \in S_{fin}(B_\Omega, B_\Omega)$.

Proof. (1) ⇒ (2). Let X be a set of reals satisfying the hypotheses and β be a countable base of X. Consider a sequence $\{\mathcal{B}_i\}_{i\in\omega}$ of countable Borel ω-covers of X where $\mathcal{B}_i = \{W_i^j\}_{j\in\omega}$ for each $i \in \omega$.

Consider a topology τ generated by the family $\mathcal{P} = \{W_i^j \cap A_s : i, j \in \omega$ and $A_s \in \beta\} \bigcup \{(X \setminus W_i^j) \cap A_s : i, j \in \omega$ and $A_s \in \beta\}$.

A space $Z = (X, \tau)$ is a separable metrizable space. Note that $\{\mathcal{B}_i\}$ is countable open ω-cover of Z for each $i \in \omega$. Let φ be a Borel mapping $\varphi : X \mapsto Z$ from a space X on a separable metric space Z. Then $B(\varphi(X)) = B(Z)$ is a dense subset of $B(X)$ and, hence, $B(Z)$ has property $S_{fin}(\mathcal{D}, \mathcal{D})$. Since $C_p(Z)$ is a dense subset of $B(Z)$, $C_p(Z)$ also has property $S_{fin}(\mathcal{D}, \mathcal{D})$.

By Theorem 21 in [2] and [9], the space Z has property $S_{fin}(\Omega, \Omega)$. It follows that there is a sequence $\{F_i = \{W_i^{j_1}, ..., W_i^{j_s}\}\}_{i\in\omega}$ such that $F_i \subset \mathcal{B}_i$ and $\bigcup_{i\in\omega} F_i \in \Omega$. It follows that $\bigcup_{i\in\omega} F_i$ is Borel ω-cover of X.

(2) ⇒ (1). Assume that X has property $S_{fin}(B_\Omega, B_\Omega)$. Let $\{D_i\}_{i\in\omega}$ be a sequence countable dense subsets of $B(X)$. We claim that for any $f \in B(X)$ there is a sequence $\{F_k\}$ of finite subsets of $B(X)$ such that $F_k \subset D_k$ for each $k \in \omega$ and $f \in \overline{\bigcup\{F_k : k \in \omega\}}$.

Without loss of generality we can assume $f = \mathbf{0}$. Denote $W_i^k = \{x \in X : -\frac{1}{k} < f_i^k(x) < \frac{1}{k}\}$ for each $f_i^k \in D_k = \{f_i^k : i \in \omega\}$ and $k \in \omega$.

If for each $j \in \omega$ there is $k(j)$ such that $W_{i(j)}^{k(j)} = X$, then a sequence $f_{k(j)} = f_{i(j)}^{k(j)}$ uniform convergence to f and, hence, $f \in \overline{\{f_{k(j)} : j \in \omega\}}$.

We can assume that $W_i^k \neq X$ for any $k, i \in \omega$.

(a). $\{W_i^k\}_{i\in\omega}$ a sequence of Borel sets of X.

(b). For each $k \in \omega$, $\{W_i^k\}_{i\in\omega}$ is a ω-cover of X.

By (2), X has property $S_{fin}(B_\Omega, B_\Omega)$, hence, there is a sequence $\{S_k = \{W_{i(1)}^k, ..., W_{i(s(k))}^k\}\}_{k\in\omega}$ such that $S_k \subset \{W_i^k\}_{i\in\omega}$ for each $k \in \omega$ and $\bigcup_{k\in\omega} S_k$ is a ω-cover of X.

Consider $\{f^k_{i(j)}\}_{j=1,...,s(k),k\in\omega}$ and we claim that $f \in \overline{\{f^k_{i(j)}\}_{j=1,...,s(k),k\in\omega}}$.

Let K be a finite subset of X, $\epsilon > 0$ and $U =< f, K, \epsilon >$ be a base neighborhood of f, then there is $k_0 \in \omega$ such that $\frac{1}{k_0} < \epsilon$ and $K \subset W^{k_0}_{i(j')}$. It follows that $f^{k_0}_{i(j')} \in U$ for some $j' \in \{1, ..., s(k_0)\}$.

Let $D = \{d_n : n \in \omega\}$ be a dense subspace of $B(X)$. Given a sequence $\{D_i\}_{i\in\omega}$ of dense subspace of $B(X)$, enumerate it as $\{D_{n,m} : n, m \in \omega\}$. For each $n \in \omega$, pick finite set $F_{n,m} \subset D_{n,m}$ so that $d_n \in \overline{\bigcup_{m\in\omega} F_{n,m}}$. Then $\bigcup_{m,n\in\omega} F_{n,m}$ is dense in $B(X)$.

□

4. Question of A. Bella, M. Bonanzinga and M. Matveev

In [3], Question 4.3, it is asked to find a sequentially separable selectively separable space which is not selective sequentially separable.

Theorem 4.1. (CH) *There is a consistent example of a space Z, such that Z is sequentially separable selectively separable, not selective sequentially separable.*

Proof. By Theorem 40 and Corollary 41 in [10], there is a $\mathfrak{c}$-Lusin set X which has property $S_1(B_\Omega, B_\Omega)$, but X does not have property $U_{fin}(\Gamma, \Gamma)$ and, hence, X does not have property $S_{fin}(B_\Gamma, B_\Gamma)$.

Consider a space $Z = B_1(X)$.

The Velichko's Theorem in [13] shows that a space $B_1(X)$ is sequentially separable for any separable metric space X.

(a). Z is sequentially separable.

By Theorem 3.3, $B(X) \in S_1(\mathcal{D}, \mathcal{D})$. Since Z is dense subset of $B(X)$ we have that $Z \in S_1(\mathcal{D}, \mathcal{D})$ and, hence, $Z \in S_{fin}(\mathcal{D}, \mathcal{D})$.

(b). Z is selectively separable.

Since X does not have property $S_{fin}(B_\Gamma, B_\Gamma)$, by Theorem 3.1, $B_1(X)$ does not have property $S_{fin}(\mathcal{S}, \mathcal{S})$.

(c). Z is not selective sequentially separable.

□

5. Question of A. Bella and C. Costantini.

In [4], Question 2.7, it is asked to find a compact T_2 sequentially separable space which is not selective sequentially separable.

Theorem 5.1. ($\mathfrak{b} < \mathfrak{q}$) *There is a consistent example of a compact T_2 sequentially separable space which is not selective sequentially separable.*

Proof. Let D be a discrete space of size $\mathfrak{b}$. Since $\mathfrak{b} < \mathfrak{q}$, a space $2^{\mathfrak{b}}$ is a sequentially separable (see Proposition 3 in [8]).

We claim that $2^{\mathfrak{b}}$ is not selective sequentially separable.

On the contrary, suppose that $2^{\mathfrak{b}}$ is selective sequentially separable. Since $non(S_{fin}(B_\Gamma, B_\Gamma)) = \mathfrak{b}$ (see Theorem 1 and Theorem 27 in [10]), there is a set of reals X such that $|X| = \mathfrak{b}$ and X has not property $S_{fin}(B_\Gamma, B_\Gamma)$. Hence there exists sequence $\{A_n : n \in \omega\}$ of elements of B_Γ that for any sequence $\{B_n : n \in \omega\}$ of finite sets such that for each n, $B_n \subseteq A_n$, we have that $\bigcup_{n\in\omega} B_n \notin B_\Gamma$.

Consider the identity mapping $id : D \mapsto X$ from the space D onto the space X. Denote $C^i_n = id^{-1}(A^i_n)$ for each $A^i_n \in A_n$ and $n, i \in \omega$. Let $C_n = \{C^i_n\}_{i\in\omega}$ and $\mathcal{S} = \{h_i\}_{i\in\omega}$ be a countable sequentially dense subset of $B(D, 2) = 2^{\mathfrak{b}}$.

For each $n \in \omega$ we consider a countable sequentially dense subset $\mathcal{S}_n$ of $B(D, 2)$ where

$\mathcal{S}_n = \{f^i_n\} := \{f^i_n \in B(D, 2) : f^i_n \restriction C^i_n = h_i$ and $f^i_n \restriction (X \setminus C^i_n) = 1$ for $i \in \omega\}$.

Since $C_n = \{C^i_n\}_{i\in\omega}$ is a Borel γ-cover of D and $\mathcal{S}$ is a countable sequentially dense subset of $B(D, 2)$, we have that $\mathcal{S}_n$ is a countable sequentially dense subset of $B(D, 2)$ for each $n \in \omega$.

Really, let $h \in B(D, 2)$, there is a sequence $\{h_s\}_{s\in\omega} \subset \mathcal{S}$ such that $\{h_s\}_{s\in\omega}$ converge to h. We claim that $\{f^s_n\}_{s\in\omega}$ converge to h. Let $K = \{x_1, ..., x_k\}$ be a finite subset of D, $\epsilon > 0$ and $W =< h, K, \epsilon >:= \{g \in B(D, 2) : |g(x_j) - h(x_j)| < \epsilon$ for $j = 1, ..., k\}$ be a base neighborhood of h, then there is a number m_0 such that $K \subset C^i_n$ for $i > m_0$ and $h_s \in W$ for $s > m_0$. Since $f^s_n \restriction K = h_s \restriction K$ for each $s > m_0$, $f^s_n \in W$ for each $s > m_0$. It follows that a sequence $\{f^s_n\}_{s\in\omega}$ converge to h.

Since $B(D, 2)$ is selective sequentially separable, there is a sequence $\{F_n = \{f^{i_1}_n, ..., f^{i_{s(n)}}_n\} : n \in \omega\}$ such that for each n, $F_n \subset \mathcal{S}_n$, and $\bigcup_{n\in\omega} F_n$ is a countable sequentially dense subset of $B(D, 2)$.

For $\mathbf{0} \in B(D, 2)$ there is a sequence $\{f^{i_{j(n)}}_{n_j}\}_{j\in\omega} \subset \bigcup_{n\in\omega} C_n$ such that $\{f^{i(n)_j}_{n_j}\}_{j\in\omega}$ converge to $\mathbf{0}$. Consider a sequence $\{C^{i_{j(n)}}_{n_j} : j \in \omega\}$.

(1) $C^{i_{j(n)}}_{n_j} \in C_{n_j}$.

(2) $\{C^{i_{j(n)}}_{n_j} : j \in \omega\}$ is a γ-cover of D.

Really, let K be a finite subset of D and $U =< \mathbf{0}\ , K, \frac{1}{2} >$ be a base neighborhood of $\mathbf{0}$, then there is a number j_0 such that $f_{n_j}^{i_{j(n)}} \in U$ for any $j > j_0$. It follows that $K \subset C_{n_j}^{i_{j(n)}}$ for any $j > j_0$. It follows that $\bigcup_{n\in\omega} C_n \in B_\Gamma$ in space D, and, hence, $\bigcup_{n\in\omega} B_n \in B_\Gamma$ in space X, a contradiction. □

Let $\mu = \min\{\kappa : 2^\kappa$ is not selective sequentially separable$\}$. It is well-known that $\mathfrak{p} \leq \mu \leq \mathfrak{q}$ (see [3]).

Theorem 5.2. $\mu = \min\{\mathfrak{b}, \mathfrak{q}\}$.

Proof. Let $\kappa < \min\{\mathfrak{b}, \mathfrak{q}\}$. Then, by Proposition 3 in [8], 2^κ is a sequentially separable space.

Let X be a set of reals such that $|X| = \kappa$ and X be a Q-set.

Analogously proof of implication (2) $\Rightarrow$ (1) in Theorem 3.1, we can claim that $B(X, 2) = 2^X = 2^\kappa$ is selective sequentially separable.

It follows that $\mu \geq \min\{\mathfrak{b}, \mathfrak{q}\}$.

Since $\mu \leq \mathfrak{q}$ we suppose that $\mu > \mathfrak{b}$ and $\mathfrak{b} < \mathfrak{q}$. Then, by Theorem 5.1, $2^\mathfrak{b}$ is not selective sequentially separable. It follows that $\mu = \min\{\mathfrak{b}, \mathfrak{q}\}$. □

In [3], Question 4.12 : is it the case $\mu \in \{\mathfrak{p}, \mathfrak{q}\}$?

By Theorem 8 in [1], there are models of set theory such that:

1. $\mu = \mathfrak{p} = \mathfrak{b} < \mathfrak{q}$;
2. $\mathfrak{p} < \mu = \mathfrak{b} = \mathfrak{q}$.

References

[1] T. Banakh, M. Machura, L. Zdomskyy, *On critical cardinalities related to Q-sets*, Mathematical Bulletin of the Shevchenko Scientific Society, 11 (2014), 21–32.

[2] A. Bella, M. Bonanzinga, M. Matveev, *Variations of selective separability*, Topology and its Applications, 156, (2009), 1241–1252.

[3] A. Bella, M. Bonanzinga, M. Matveev, *Sequential+separable vs sequentially separable and another variation on selective separability*, Cent. Eur. J. Math. 11-3 (2013), 530–538.

[4] A. Bella, C. Costantini, *Sequential Separability vs Selective Sequential Separability*, Filomat 29:1, (2015), 121–124.

[5] L. Bukovský, J. Šupina, *Sequence selection principles for quasi-normal convergence*, Topology and its Applications, 159, (2012), p.283–289.

[6] E.K. van Douwen, *The integers and topology*, in: Handbook of Set-Theoretic Topology, North-Holland, Amsterdam, (1984).

[7] A.V. Osipov, E.G. Pytkeev, *On sequential separability of functional spaces*, Topology and its Applications, (2017).

[8] P. Gartside, J.T.H. Lo, A. Marsh, *Sequential density*, Topology and its Applications, 130, (2003), p.75–86.

[9] W. Just, A.W. Miller, M. Scheepers, P.J. Szeptycki, *The combinatorics of open covers, II*, Topology and its Applications, 73, (1996), 241–266.

[10] M. Scheepers, B. Tsaban, *The combinatorics of Borel covers*, Topology and its Applications, 121, (2002), p.357–382.

[11] M. Scheepers, *Combinatorics of open covers VI: Selectors for sequences of dense sets*, Quaestiones Mathematicae, 22, (1999), p. 109–130.

[12] B. Tsaban, L. Zdomskyy, *Hereditarily Hurewicz spaces and Arhangel'skiĭ sheaf amalgamations*, Journal of the European Mathematical Society, 12, (2012), 353–372.

[13] N.V. Velichko, *On sequential separability*, Mathematical Notes, Vol.78, Issue 5, 2005, p. 610–614.